

Analysis at the Sequence Level of Mutations Induced by the Ultimate Carcinogen *N*-Acetoxy-*N*-2-Acetylaminofluorene

by Robert P. P. Fuchs,* Nicole Schwartz*
and Michel P. Daune*

The covalent binding of an ultimate carcinogen to the DNA bases or phosphate groups creates a premutational lesion that *in vivo* is processed by the repair, replication and recombination enzymes, and eventually may be converted into a mutation. Being interested in the way that an initial premutational event is converted into a stable heritable mutation, we have sequenced stable mutations in a gene that has formed covalent adducts *in vitro* with *N*-acetoxy-*N*-2-acetylaminofluorene (*N*-AcO-AAF), a model for the ultimate metabolite of the rat liver carcinogen 2-acetylaminofluorene, AAF). *In vivo* studies have shown the mutagenicity of AAF and its derivatives in both bacterial and eukaryotic systems. *N*-AcO-AAF reacts *in vitro* with DNA leading mainly to the formation of a guanine adduct, *N*-2-(deoxyguanosin-8-yl)-acetylaminofluorene (80%) and to at least three minor adducts. Studies by our group showed that binding of *N*-AcO-AAF to DNA resulted in a local distortion of the DNA helix around the C-8 adduct (the insertion-denaturation model).

We describe here the analysis of forward mutations induced in the tetracycline-resistance gene of pBR322 by directing the chemical reaction of the carcinogen to a small restriction fragment (*Bam*HI-*Sal*I) inside the antibiotic-resistance gene. Mutants are selected for ampicillin (Ap) resistance and tetracycline (Tc) sensitivity. The plasmid DNA of such mutants was analyzed for sequence changes in the fragment where the AAF binding had been directed.

We show here that the mutations are mainly frameshifts involving GC base pairs and that certain base pairs (hotspots) are affected at high frequencies.

Introduction

An important step in the carcinogenic process is thought to be the initial attack of the DNA molecule by a so-called ultimate carcinogen. In fact, more than 90% of the carcinogens tested are mutagens in bacterial systems (1). The premutational event is the covalent binding of the ultimate carcinogen to the DNA bases or phosphate groups. The chemical structure of the adducts formed, and to a lesser extent the structural changes induced in the DNA double helix in the neighborhood of the adducts, has been extensively studied during the last ten years. However the crucial question is "How will the different repair, replication and recombination enzymes handle these chemically modified bases?" In other words, since the end point of this initial step is a mutation, "How is this initial premutational event converted into a stable and heritable

mutation?" *N*-Acetoxy-*N*-2-acetylaminofluorene (*N*-AcO-AAF) is a model ultimate metabolite of the strong rat liver carcinogen 2-acetylaminofluorene (AAF). *In vivo* studies have shown the mutagenicity of AAF and its derivatives in both bacterial (2, 3) and eukaryotic systems (4). *N*-AcO-AAF reacts *in vitro* with DNA leading mainly to the formation of a guanine adduct (5), *N*-2-(deoxyguanosin-8-yl)-acetylaminofluorene (80%) and also to at least three minor adducts (N. Schwartz, R. P. P. Fuchs and M. P. Daune, unpublished results), one of which is characterized as 3-(deoxyguanosin-*N*²-yl)-acetylaminofluorene (6).

Studies from our group led to the general conclusion that binding of *N*-AcO-AAF to DNA resulted in a local distortion of the DNA helix around the C-8 adduct (7-9). We have called this structural alteration the insertion-denaturation model (10). A similar model has been proposed by other investigators (11).

In this paper we describe the analysis of forward mutations induced in the tetracycline-resistance

*Institut de Biologie Moléculaire et Cellulaire du CNRS, 15 rue Rene Descartes, 67084 Strasbourg Cédex, France

The sequences appearing in this table are the wild type sequences with the numbering defined by Sutcliffe (20). The bases involved in the deletion mutations are boxed with dotted lines. The different possibilities to obtain a given mutated sequence are shown. Mutant 36 exhibits two mutations: a - 2 deletion as in mutant 34 and a + 1 addition of a C residue within the sequence CCC at positions 526-528. Mutant 35 has got a - 1 deletion of a G at position 416 and a double transition, GC \rightarrow AT, at position 414-415.




FIGURE 3. Part of the sequence of mutant 34 and of the wild type DNA showing the -2 deletion of a CG doublet within the hot spot sequence GGCGCC.

guanine-AAF adduct that might favor the hairpin structure shown in Figure 4. Due to the multicopy state and to the recessivity of the mutations that are scored in our system, the conversion of the premutagenic lesion into a stable mutation most likely occurs simultaneously in both strands prior to replication.




FIGURE 4. Hypothetical hairpin structure at hot spot sequences 1 and 2. According to the insertion-denaturation model proposed by Fuchs and co-workers (7, 10), there is a local denaturation of the helix around the guanine-AAF adduct that might favor the hairpin structure.

The molecular mechanism by which the mutation is being fixed remains to be elucidated.

This work has been supported by Grant No. 79.7.0664 from the D.G.R.S.T. (Délégation Générale à la Recherche Scientifique et Technique).

REFERENCES

- McCann, J., Choi, E., Yamasaki, E., and Ames, B. N. Detection of carcinogens as mutagens in the Salmonella/microsome test: Assay of 300 carcinogens. *Proc. Natl. Acad. Sci. (U.S.)* 72: 5135-5139 (1975).
- Ames, B. N., Gurney, E. G., Miller, J. A., and Bartsch, H. Carcinogens as frameshift mutagens: metabolites and derivatives of 2-acetylaminofluorene and other aromatic amine carcinogens. *Proc. Natl. Acad. Sci. (U.S.)* 70: 782-786 (1973).
- Santella, R. M., Fuchs, R. P. P., and Grunberger, D. Mutagenicity of 7-iodo and 7-fluoro derivatives of *N*-hydroxy and *N*-acetoxy-*N*-2-acetylaminofluorene in the *Salmonella typhimurium* assay. *Mutat. Res.* 67: 85-87 (1979).
- Landolph, J. R., and Heidelberger, C. Chemical carcinogens produce mutations to ouabain resistance in transformable C 3H/10 T $\frac{1}{2}$ Cl 8 mouse fibroblasts. *Proc. Natl. Acad. Sci. (U.S.)* 76: 930-934 (1979).
- Kriek, E., Miller, J. A., Juhl, V., and Miller, E. C. 8-(*N*-2-Fluorenylacetyl)guanosine and arylamidation reaction product of guanosine and the carcinogen *N*-acetoxy-*N*-2-fluorenylacetylamine in neutral solution. *Biochemistry* 6: 177-182 (1967).
- Westra, J. G., Kriek, E., and Hittenhausen, H. Identification of the persistently bound form of the carcinogen *N*-acetyl-2-aminofluorene to rat liver DNA *in vivo*. *Chem.-Biol. Interact.* 15: 149-164 (1976).
- Fuchs, R., and Daune, M. Physical studies on deoxyribonucleic acid after covalent binding of a carcinogen. *Biochemistry* 11: 2659-2666 (1972).
- Fuchs, R. P. P., and Daune, M. P. Dynamic structure of DNA modified with the carcinogen *N*-acetoxy-*N*-2-acetylaminofluorene. *Biochemistry* 13: 4435-4440 (1974).
- Fuchs, R. P. P. *In vitro* recognition of carcinogen-induced local denaturation sites in native DNA by S1 endonuclease from *Aspergillus oryzae*. *Nature*, 257: 151-152 (1975).
- Fuchs, R. P. P., Lefèvre, J. F., Pouyet, J., and Daune, M. P. Comparative orientation of the fluorene residues in native DNA modified by *N*-acetoxy-*N*-2-acetylaminofluorene and two 7-halogen derivatives. *Biochemistry* 15: 3347-3351 (1976).
- Grunberger, D., and Weinstein, I. B. Conformational changes in nucleic acids modified by chemical carcinogens. In: *Chemical Carcinogens and DNA* (P. Grover, Ed.), CRC Press, Vol. 2, Boca Raton, FL, pp. 59-94.
- Fuchs, R. P. P., Schwartz, N., and Daune, M. P. Hot spots of frameshift mutations induced by the ultimate carcinogen *N*-acetoxy-*N*-2-acetylaminofluorene. *Nature* 294: 657-659 (1981).
- Howard-Flanders, P., Boyce, R. P., and Theriot, L. Three loci in *E. coli* K-12 that control the excision of pyrimidine dimers and certain other mutagen products from DNA. *Genetics* 53: 1119-1136 (1966).
- Lefèvre, J. F., Fuchs, R. P. P., and Daune, M. P. Comparative studies on the 7-iodo and 7-fluoro derivatives of *N*-acetoxy-*N*-2-acetylaminofluorene: binding sites on DNA and conformational change of modified deoxytrinucleotides. *Biochemistry* 17: 2561-2567 (1978).
- de Murcia, G., Lang, M. C., Freund, A. M., Fuchs, R. P. P., Daune, M. P., Sage, E., and Leng, M. Electron microscopic visualization of *N*-acetoxy-*N*-2-acetylaminofluorene binding sites in Col E1 DNA by means of specific antibodies. *Proc. Natl. Acad. Sci. (U.S.)* 76: 6076-6080 (1979).
- Cohen, S. N., Chang, A. C. Y., and Hsu, L. Non-chromosomal antibiotic resistance in bacteria: genetic transformation of *Escherichia coli* by R-factor DNA. *Proc. Natl. Acad. Sci. (U.S.)* 69: 2110-2114 (1972).
- Clewell, D. B., and Helinski, D. R. Supercoiled circular DNA-protein complex in *Escherichia coli*: purification and induced conversion to an open circular DNA form. *Proc. Natl. Acad. Sci. (U.S.)* 62: 1159-1166 (1969).
- Katz, L., Kingsbury, D. K., and Helinski, D. R. Stimulation by cyclic adenosine monophosphate of plasmid deoxyribonucleic acid-protein relaxation complex. *J. Bacteriol.* 114: 557-591 (1973).
- Maxam, A. M., and Gilbert, W. A new method for sequencing DNA. *Proc. Natl. Acad. Sci. (U.S.)* 74: 560-564 (1977).
- Sutcliffe, J. G. Complete nucleotide sequence of the *Escherichia coli* plasmid pBR 322. *Cold Spring Harbor Symp. Quant. Biol.* 43: 77-90 (1979).
- Isono, K., and Yournon, J. Chemical carcinogens as frameshift mutagens: *Salmonella* DNA sequence sensitive to mutagenesis by polycyclic carcinogens. *Proc. Natl. Acad. Sci. (U.S.)* 71: 1612-1617 (1974).